0 13013 54210 8

PARKER'S

Super SOILIFE

WEED & FEED 10-6-4

TRIKOTE TECHNOLOGY INSIDE

PARKER'S®

QUALITY FERTILIZERS SINCE 1904

Super SOILIFE®

WEED & FEED

HERBICIDA Y ALIMENTO

10-6-4

- Kills dandelions and other broadleaf weeds
  Elimina el diente de león y muchas otras malezas de hoja ancha
- Promotes a green lawn and strong roots
  Favorece el desarrollo de césped verde y raíces fuertes

Timed-Release Nitrogen for Extended Feeding

Nitrógeno de liberación lenta para una nutrición prolongada

TRIKOTE® TECHNOLOGY INSIDE

ACTIVE INGREDIENTS:

2,4-D	0.64%
Mecoprop-p	0.16%
Dicamba	0.03%
OTHER INGREDIENTS:	99.17%
TOTAL	100.00%

CAS Registry Numbers:
2,4-D CAS# 94-75-7
Mecoprop-p CAS# 16484-77-8,
Dicamba CAS# 1918-00-9

THIS PRODUCT CONTAINS:
12.80 lbs 2,4-dichlorophenoxyacetic acid equivalent per ton.
3.20 lbs (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
0.60 lbs dicamba (3,6-dichloro-o-anisic acid) equivalent per ton.
Isomer Specific by AOAC Methods
TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional Precautionary Statements & First Aid.
MANTENGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCION Vea el panel posterior para conocer las precauciones adicionales y los primeros auxilios.

Net Wt 16 lb (7.2 kg)

PARKER'S Super SOILIFE WEED & FEED 10-6-4 TRIKOTE TECHNOLOGY INSIDE

0 13013 54210 8

PARKER'S
QUALITY FERTILIZERS SINCE 1964

Super SOILIFE®

WEED & FEED 10-6-4

COVERS UP TO 5,000 SQ FT (50 ft x 100 ft)

GUARANTEED ANALYSIS 10-6-4

Total Nitrogen (N)	10%
2.3 % Ammoniacal Nitrogen	
7.7 % Urea Nitrogen*	
Available Phosphate (P_2O_5)	6%
Soluble Potash (K_2O)	4%

Derived from: Polymer-coated sulfur-coated urea, urea, ammonium phosphate and muriate of potash.

*1.5% Slowly available nitrogen from coated urea.

F1462

GENERAL TIPS

1. Determine size of area to be treated – measure length and width in feet, then multiply to determine coverage area in sq ft (length x width). Next, subtract non-lawn areas.
2. Use the spreader setting chart to determine the appropriate setting.
3. Treat as directed – thorough coverage is important.

READ THE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE

This product controls a wide range of broadleaf weeds in home lawns. In addition, the fertilizer provides a supply of plant food.

FOR USE ON

Bahiagrass
Bermudagrass
Buffalograss
Centipedegrass
Fescue
Kentucky bluegrass
Perennial ryegrass
St. Augustinegrass
Zoysiagrass

USE PRECAUTIONS

- Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.
- Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental shrubbery.
- Do not exceed the application rate for any area. When applying under trees in an open lawn, walk past trees in a straight path. Circling trees will cause you to pass over some areas twice so that too much material is applied.
- If overseeding wait 4 weeks after grass seeds sprout before applying this product.

APPLICATION INSTRUCTIONS

STEP 1 Mow lawn to normal height 1 to 2 days before application.
STEP 2 Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
STEP 3 Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
STEP 4 Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

WEEDS CONTROLLED/MALEZAS CONTROLADAS

Annual yellow sweetclover
Aster
Austrian fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, field
Bird vetch
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Black mustard
Blackseed plantain
Blessed thistle
Bloodflower milkweed
Blue lettuce
Blue vervain
Bracted plantain
Bristly oxtongue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn
Buckhorn plantain
Bulbous buttercup
Bull thistle
Bullnettle
Burclover
Burdock
Burning nettle
Burweed
Buttercup
Buttonweed
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catnip
Catsear
Chickweed, common
Chickweed, mouseear
Chicory
Cinquefoil
Clover, crimson
Clover, hop
Clover, red
Clover, strawberry
Clover, sweet
Clover, white
Cockle
Cocklebur
Common mullein
Creeping jenny
Cudweed
Curly dock
Daisy, English
Daisy fleabane
Daisy, oxeye
Dandelion
Dichondra
Dogfennel
Dollarweed
False dandelion
Falseflax
False sunflower
Fiddleneck
Florida pusley
Frenchweed
Galinsoga
Goldenrod
Ground ivy
Gumweed
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Heath aster
Hedge bindweed
Hedge mustard
Hemp
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Horsenettle
Jimsonweed
Knawel
Knotweed
Kochia
Lambsquarters
Lespedeza
Mallow
Matchweed
Mexicanweed
Milk vetch
Morningglory
Mouseear hawkweed
Mugwort
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oriental cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis
Plantain
Poison ivy
Poison oak
Pokeweed
Poorjoe
Prairie sunflower
Prickly lettuce
Prickly sida
Prostrate knotweed
Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncturevine
Purslane, common
Ragweed
Red sorrel
Redroot pigweed
Redstem filaree
Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Spotted spurge
Spurweed
Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy ragwort
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild mustard
Wild marigold
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower
pepperweed and other broadleaf weeds

HOW MUCH TO APPLY

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass:

Apply 3.2 lb of product per 1,000 sq ft of turf grass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 lb of product per 1,000 sq ft should be used 30 days after the first application.

Do not apply this product to — 'Floratam' St. Augustinegrass in Florida.

The maximum application rate to turf is 1.1 lb 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

A lawn typically utilizes 1 to 4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.32 lb of nitrogen per 1,000 sq ft. Consult your local Extension service for questions regarding a fertilizer program suitable to your geographical region.

SPREADERS AND CALIBRATION

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

Single Pass — Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.

IMPORTANT • It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult owners' manual. ***NEVER APPLY BY HAND***

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.

Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers 5,000 sq ft.

SPREADER SETTINGS

Rotary Spreaders	
Scotts Edge Guard	6.75
Scotts SpeedyGreen	6.75
Earthway 2050	9.0
Drop Spreader	
Scotts AccuGreen	7.5

CALIBRATION INSTRUCTIONS

- For 3.2 lb of product per 1,000 sq ft of turfgrass: Check your spreader on 310 sq ft and adjust to apply at the rate of 1 lb of product per 310 sq ft.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.
CONTAINER DISPOSAL:
If empty: Do not reuse this container. Place in trash or offer for recycling if available.
If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. When using this product, wear long-sleeved shirt, long pants, socks and shoes.
After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-874-8892 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

CONSEJOS GENERALES

1. Calcule el área que se va a tratar; mida el largo y el ancho en pies y después multiplique para calcular el área de cobertura en pies cuadrados (largo x ancho). Luego reste las áreas que no tengan césped.
2. Use el cuadro de ajustes del esparcidor para verificar el ajuste apropiado.
3. Realice el tratamiento como se indica; es importante que la cobertura sea total.

ALTO! LEA PRIMERO LA ETIQUETA. CUMPLA TODAS LAS PRECAUCIONES Y SIGA CUIDADOSAMENTE LAS INSTRUCCIONES.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

No debe haber personas (salvo quien realice la aplicación) ni mascotas en el área por tratar durante la aplicación. No entre en las áreas de tratamiento hasta que el polvo se haya asentado.

No se debe usar en césped que se cultive para la venta u otro uso comercial, como panes de césped, ni para la producción de semillas comerciales, ni con fines de investigación.

DONDE UTILIZARLO

Este producto controla una gran variedad de malezas de hoja ancha en céspedes residenciales. Además, el fertilizante proporciona nutrientes para las plantas.

PARA USAR EN

Bahía
Bermuda
Búfalo
Festuca
Césped azul de Kentucky
Césped ballico perenne
Zoysia

PRECAUCIONES SOBRE EL MODO DE EMPLEO

- No aplique este producto en césped alfombra, dichondra, plantas ornamentales ni céspedes que contengan tréboles deseables.
- Escoja un día en que no haya viento para asegurar una cobertura uniforme y evitar que los gránulos se diseminen sobre flores, hortalizas y arbustos ornamentales.
- No aplique una dosis mayor a la recomendada en ninguna zona. Cuando lo aplique debajo de los árboles en un césped abierto, camine en línea recta al lado de los árboles. Rodear los árboles en círculo lo hará pasar dos veces por algunas zonas y por lo tanto aplicar demasiado producto.
- Si sobresiembra, espere 4 semanas después de que las semillas de césped germinen para aplicar este producto.

INSTRUCCIONES DE APLICACION

PASO 1 Corte el césped hasta la altura normal 1 a 2 días antes de la aplicación.
PASO 2 Riegue muy bien el césped por lo menos 1 día antes de la aplicación para mantener la humedad hasta el siguiente riego (vea el paso 4).
PASO 3 Aplique el producto cuando las malezas son jóvenes y se encuentran en la etapa de crecimiento activo, de preferencia por la mañana, cuando hay rocío sobre el pasto. Si el pasto no está húmedo en el momento de la aplicación, rocíelo ligeramente con agua para que los gránulos se adhieran y permanezcan sobre la superficie de las hojas de las malezas.
PASO 4 No riegue ni rocíe durante 1 a 2 días después de la aplicación. Riegue, rocíe o moje completamente el césped a los 2 a 3 días de la aplicación.

CANTIDAD POR APLICAR

Para césped azul de Kentucky, festucas, césped ballico perenne, césped Bermuda, Búfalo, Bahía, Zoysia, San Agustín y ciempiés:

Aplique 3.2 libras de producto cada 1,000 pies cuadrados de césped. Puede ser necesaria una segunda aplicación o aplicación de control en caso de infestaciones densas de malezas o debido a condiciones estacionales desfavorables. La segunda aplicación de 3.2 libras de producto cada 1,000 pies cuadrados se debe hacer a los 30 días de la primera aplicación.

No aplique este producto a la variedad 'Floratam' del césped San Agustín en Florida.

La dosis máxima de aplicación al césped es de 1.1 libra de equivalente ácido de 2,4-D por acre, por aplicación y por sitio. La cantidad máxima de aplicaciones a voleo por lugar de tratamiento es de 2 por año.

Es esencial que el producto se distribuya uniformemente en el área tratada. Escoja un día en que no haya viento para asegurar una cobertura uniforme y evitar que los gránulos se diseminen sobre flores, hortalizas y arbustos ornamentales. No aplique una dosis mayor a la recomendada en ninguna zona y tenga cuidado cuando aplique este producto dentro de la franja de goteo de árboles, arbustos y otras plantas ornamentales.

En general, un césped usa entre 1 y 4 libras de nitrógeno cada 1,000 pies cuadrados por temporada de crecimiento. Cuando se aplica según las instrucciones, este producto proporciona 0.32 libras de nitrógeno cada 1,000 pies cuadrados. Consulte a la oficina local del servicio de extensión para obtener información sobre un programa de aplicación de fertilizante adecuado para su región geográfica.

ESPARCIDORES Y CALIBRACION

Los esparcidores por goteo y los esparcidores giratorios se deben calibrar para lograr la aplicación uniforme de este producto.

Paso único — Opere el equipo a intervalos correspondientes al ancho o a la banda de aspersión del esparcidor. Con esparcidores por goteo, superponga los carriles para asegurar la cobertura total. Cierre las aberturas del esparcidor al girar para cambiar de dirección o cuando se detenga.

IMPORTANTE • Es importante seguir las instrucciones tal como se indican en este producto de calidad para el cuidado del césped, ya que aplicar un exceso de producto puede causar daños a su césped. Los ajustes del esparcidor son aproximados. Puede haber diferencias como consecuencia del estado del esparcidor, la velocidad a la cual se aplica y la irregularidad de la zona tratada. Asegúrese de revisar el esparcidor antes de comenzar y de ajustarlo si es necesario. Consulte el manual del propietario. ***NUNCA LO APLIQUE A MANO.***

Para evitar que queden áreas y líneas sin producto, primero esparza sobre el borde del césped y después complete el centro aplicando en forma paralela y uniforme. Camine a paso regular para aplicar uniformemente el producto en todo el césped.

Cobertura: Para calcular cuánto necesita aplicar, multiplique el largo de su césped por el ancho. Esto equivale a la superficie total en pies cuadrados. Reste todas las áreas que no sean de césped (casa, jardín, etc.). Una bolsa alcanza para cubrir 5,000 pies cuadrados.

AJUSTES DEL ESPARCIDOR

Esparcidores giratorios	
Scotts Edge Guard	6.75
Scotts SpeedyGreen	6.75
Earthway 2050	9.0
Esparcidor por goteo	
Scotts AccuGreen	7.5

INSTRUCCIONES DE CALIBRACION

- Para 3.2 libras de producto cada 1,000 pies cuadrados de césped: verifique que su esparcidor esté regulado en 310 pies cuadrados y ajuste para aplicar la dosis de 1 libra de producto cada 310 pies cuadrados.

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO: Guarde el producto en su envase original y conserve todas las bolsas que contengan producto sin usar en un lugar cerrado, inaccesible a los niños y a los animales domésticos
ELIMINACION DEL ENVASE:
Si está vacío: No vuelva a usar el envase. Tírelo en la basura, o puede ofrecerlo para reciclarlo, si existe esa opción. **Si está parcialmente vacío:** Llame al organismo local que se encarga de los desechos sólidos o llame gratis al 1-800-CLEANUP (1-800-253-2687) para obtener instrucciones sobre su eliminación. Nunca arroje el producto sin usar por ningún desagüe interior ni exterior.

PRECAUCIONES

Es nocivo para los seres humanos y los animales domésticos

PRECAUCION. Es nocivo si se ingiere o si se absorbe por la piel. Causa irritación moderada a los ojos. Evite el contacto con la piel, los ojos o la ropa. Evite respirar el polvo. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines y zapatos. Después de usar este producto, quítese la ropa y lávela por separado antes de volver a usarla y rápidamente lávese muy bien las manos y la piel que estuvo expuesta, con agua y jabón.

RIESGOS PARA EL MEDIO AMBIENTE

El arrastre o el escurrimiento pueden afectar desfavorablemente a la vegetación que no sea el objetivo del tratamiento. No lo aplique directamente al agua. No contamine el agua cuando elimine las aguas de lavado del equipo.

Cuando se aplica según las instrucciones, este producto cumple con las normas para metales aceptadas por la Association of American Plant Food Control Officials.

Se puede obtener información sobre el contenido y las concentraciones de metales de este producto en la siguiente dirección de Internet: http://www.regulatory-info-sb.com

Si usted aplica este producto según las instrucciones y no queda totalmente satisfecho, envíe la prueba de compra de este paquete y el recibo original de la caja registradora a la dirección que se encuentra a continuación. Le devolveremos el valor de compra que pagó.

GARANTIA LIMITADA Y DECLINACION DE RESPONSABILIDAD.

El distribuidor solamente garantiza que la composición química de este producto concuerda con la declaración de ingredientes expresada en la etiqueta, y que el producto es razonablemente adecuado para el uso indicado cuando se aplica según el Modo de empleo.

EL DISTRIBUIDOR NO ESTABLECE NI PROPONE NINGUNA OTRA GARANTIA EXPRESA NI IMPLICITA, INCLUIDA CUALQUIER GARANTIA DE COMERCIABILIDAD O CONVENIENCIA PARA UN PROPOSITO PARTICULAR, LAS QUE SE DECLINAN EXPRESAMENTE. Esta garantía limitada no se extiende al uso del producto de manera incompatible con las instrucciones expresadas en la etiqueta, las advertencias o las precauciones, ni al uso del producto en condiciones anormales, como sequía, lluvias excesivas, tornados, huracanes, etc. Estos factores están fuera del control del distribuidor o del vendedor. Cualquier tipo de daño que surja como consecuencia del incumplimiento de la garantía del distribuidor estará limitado a los daños directos, y no incluirá daños indirectos ni emergentes tales como pérdida de utilidades ni valores, salvo lo dispuesto en otro lugar por la ley.

Los términos de esta garantía limitada y declinación de responsabilidad no se pueden modificar por ningún tipo de declaración ni acuerdo escrito ni verbal. Ningún empleado ni representante del vendedor está autorizado a modificar ni abusar de manera alguna de los términos de esta garantía limitada y declinación de responsabilidad.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-sb.com.

If you apply this product as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address below. We will refund the purchase price you paid.

LIMITED WARRANTY AND DISCLAIMER. The distributor warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE DISTRIBUTOR NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the distributor or the seller. Any damages arising from a breach of the distributor's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of the Limited Warranty and Disclaimer in any manner.

Questions or Comments? Call 1-800-874-8892.

Distributed and guaranteed by Chemsico, Division of United Industries Corporation, PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 2217-819-9688

EPA Est. Nos. 9198-OH-1(M), 9198-OH-2(B), 9198-AL-1(A), 49991-TX-1(AZ), 49991-TX-2(BC), 45065-MN-1(BD), 59144-GA-1(BE), 6665-NY-1(BF), 35512-FL-2(BG), 8378-IN-1(AS), 43854-MO-1(BH), 49710-TX-1(AT), 49710-TX-3(AU), 3234-UT-1(AV), 3234-AZ-1(AW), 8660-AL-1(BN), 8660-OH-1(BO), 74764-CA-1(BK), 74764-FL-1(BL), 9404-FL-1(AX), 35497-OR-1(AY)

Circled letter is first letter of lot number. 11-7488 **© 2005**

Trikote® is a registered trademark of RLC Technologies, L.L.C. The Trikote design is a trademark of RLC Technologies, L.L.C.

FPO
0 13013 54210 8

0 13013 54210 8

PARKER'S®

QUALITY FERTILIZERS SINCE 1904

Super SOILIFE®

WEED & FEED

PARKER'S®
Super SOILIFE®
WEED & FEED

TRIKOTE® TECHNOLOGY INSIDE

10-6-4

WEED & FEED

HERBICIDA Y ALIMENTO

10-6-4

- Kills dandelions and other broadleaf weeds
  Elimina el diente de león y muchas otras malezas de hoja ancha
- Promotes a green lawn and strong roots
  Favorece el desarrollo de césped verde y raíces fuertes

Timed-Release Nitrogen for Extended Feeding

Nitrógeno de liberación lenta para una nutrición prolongada

TRIKOTE® TECHNOLOGY INSIDE

ACTIVE INGREDIENTS:

2,4-D	0.64%
Mecoprop-p	0.16%
Dicamba	0.03%
OTHER INGREDIENTS:	99.17%
TOTAL	100.00%

CAS Registry Numbers:
2,4-D CAS# 94-75-7
Mecoprop-p CAS# 16484-77-8,
Dicamba CAS# 1918-00-9

THIS PRODUCT CONTAINS:
12.80 lbs 2,4-dichlorophenoxyacetic acid equivalent per ton.
3.20 lbs (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
0.60 lbs dicamba (3,6-dichloro-o-anisic acid) equivalent per ton.
Isomer Specific by AOAC Methods
TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional Precautionary Statements & First Aid.
MANTENGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCION Vea el panel posterior para conocer las precauciones adicionales y los primeros auxilios.

Net Wt 16 lb (7.2 kg)

PARKER'S® Super SOILIFE® WEED & FEED 10-6-4 TRIKOTE® TECHNOLOGY INSIDE

0 13013 54210 8

PARKER'S
QUALITY FERTILIZERS SINCE 1904

Super SOILIFE®

WEED & FEED 10-6-4

COVERS UP TO 5,000 SQ FT (50 ft x 100 ft)

GUARANTEED ANALYSIS 10-6-4

Total Nitrogen (N)	10%
2.3 % Ammoniacal Nitrogen	
7.7 % Urea Nitrogen*	
Available Phosphate (P_2O_5)	6%
Soluble Potash (K_2O)	4%

Derived from: Polymer-coated sulfur-coated urea, urea, ammonium phosphate and muriate of potash.

*1.5% Slowly available nitrogen from coated urea.

F1462

GENERAL TIPS

1. Determine size of area to be treated – measure length and width in feet, then multiply to determine coverage area in sq ft (length x width). Next, subtract non-lawn areas.
2. Use the spreader setting chart to determine the appropriate setting.
3. Treat as directed – thorough coverage is important.

READ THE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHERE TO USE

This product controls a wide range of broadleaf weeds in home lawns. In addition, the fertilizer provides a supply of plant food.

FOR USE ON

Bahiagrass
Bermudagrass
Buffalograss
Centipedegrass
Fescue
Kentucky bluegrass
Perennial ryegrass
St. Augustinegrass
Zoysiagrass

USE PRECAUTIONS

- Do not apply this product to carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers.
- Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables and ornamental shrubbery.
- Do not exceed the application rate for any area. When applying under trees in an open lawn, walk past trees in a straight path. Circling trees will cause you to pass over some areas twice so that too much material is applied.
- If overseeding wait 4 weeks after grass seeds sprout before applying this product.

APPLICATION INSTRUCTIONS

STEP 1 Mow lawn to normal height 1 to 2 days before application.
STEP 2 Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
STEP 3 Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
STEP 4 Do not irrigate or sprinkle for 1 to 2 days after application. Irrigate, sprinkle, or water the lawn thoroughly at 2 to 3 days after application.

WEEDS CONTROLLED/MALEZAS CONTROLADAS

Annual yellow sweetclover
Aster
Austrian fieldcress
Bedstraw
Beggarticks
Betony, Florida
Bindweed, field
Bird vetch
Bitter wintercress
Bittercress, hairy
Bitterweed
Black-eyed Susan
Black medic
Black mustard
Blackseed plantain
Blessed thistle
Buttercup
Buttonweed
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catnip
Catsear
Chickweed, common
Chickweed, mouseear
Chicory
Cinquefoil
Clover, crimson
Clover, hop
Clover, red
Clover, strawberry
Clover, sweet
Fiddleneck
Florida pusley
Frenchweed
Galinsoga
Goldenrod
Ground ivy
Gumweed
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Heath aster
Hedge bindweed
Hedge mustard
Hemp
Henbit
Hoary cress
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oriental cocklebur
Oxalis
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis
Plantain
Poison ivy
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild mustard
Wild marigold

CONSEJOS GENERALES

1. Calcule el área que se va a tratar; mida el largo y el ancho en pies y después multiplique para calcular el área de cobertura en pies cuadrados (largo x ancho). Luego reste las áreas que no tengan césped.
2. Use el cuadro de ajustes del esparcidor para verificar el ajuste apropiado.
3. Realice el tratamiento como se indica; es importante que la cobertura sea total.

ALTO! LEA PRIMERO LA ETIQUETA. CUMPLA TODAS LAS PRECAUCIONES Y SIGA CUIDADOSAMENTE LAS INSTRUCCIONES.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

No debe haber personas (salvo quien realice la aplicación) ni mascotas en el área por tratar durante la aplicación. No entre en las áreas de tratamiento hasta que el polvo se haya asentado.

No se debe usar en césped que se cultive para la venta u otro uso comercial, como panes de césped, ni para la producción de semillas comerciales, ni con fines de investigación.

DONDE UTILIZARLO

Este producto controla una gran variedad de malezas de hoja ancha en céspedes residenciales. Además, el fertilizante proporciona nutrientes para las plantas.

PARA USAR EN

Bahía
Bermuda
Búfalo
Festuca
Césped azul de Kentucky
Césped ballico perenne
Zoysia

PRECAUCIONES SOBRE EL MODO DE EMPLEO

- No aplique este producto en césped alfombra, dichondra, plantas ornamentales ni céspedes que contengan tréboles deseables.
- Escoja un día en que no haya viento para asegurar una cobertura uniforme y evitar que los gránulos se diseminen sobre flores, hortalizas y arbustos ornamentales.
- No aplique una dosis mayor a la recomendada en ninguna zona. Cuando lo aplique debajo de los árboles en un césped abierto, camine en línea recta al lado de los árboles. Rodear los árboles en círculo lo hará pasar dos veces por algunas zonas y por lo tanto aplicar demasiado producto.
- Si sobresiembra, espere 4 semanas después de que las semillas de césped germinen para aplicar este producto.

INSTRUCCIONES DE APLICACION

PASO 1 Corte el césped hasta la altura normal 1 a 2 días antes de la aplicación.
PASO 2 Riegue muy bien el césped por lo menos 1 día antes de la aplicación para mantener la humedad hasta el siguiente riego (vea el paso 4).
PASO 3 Aplique el producto cuando las malezas son jóvenes y se encuentran en la etapa de crecimiento activo, de preferencia por la mañana, cuando hay rocío sobre el pasto. Si el pasto no está húmedo en el momento de la aplicación, rocíelo ligeramente con agua para que los gránulos se adhieran y permanezcan sobre la superficie de las hojas de las malezas.
PASO 4 No riegue ni rocíe durante 1 a 2 días después de la aplicación. Riegue, rocíe o moje completamente el césped a los 2 a 3 días de la aplicación.

CANTIDAD POR APLICAR

Para césped azul de Kentucky, festucas, césped ballico perenne, césped Bermuda, Búfalo, Bahía, Zoysia, San Agustín y ciempiés:

Aplique 3.2 libras de producto cada 1,000 pies cuadrados de césped. Puede ser necesaria una segunda aplicación o aplicación de control en caso de infestaciones densas de malezas o debido a condiciones estacionales desfavorables. La segunda aplicación de 3.2 libras de producto cada 1,000 pies cuadrados se debe hacer a los 30 días de la primera aplicación.

No aplique este producto a la variedad 'Floratam' del césped San Agustín en Florida.

La dosis máxima de aplicación al césped es de 1.1 libra de equivalente ácido de 2,4-D por acre, por aplicación y por sitio. La cantidad máxima de aplicaciones a voleo por lugar de tratamiento es de 2 por año.

Es esencial que el producto se distribuya uniformemente en el área tratada. Escoja un día en que no haya viento para asegurar una cobertura uniforme y evitar que los gránulos se diseminen sobre flores, hortalizas y arbustos ornamentales. No aplique una dosis mayor a la recomendada en ninguna zona y tenga cuidado cuando aplique este producto dentro de la franja de goteo de árboles, arbustos y otras plantas ornamentales.

En general, un césped usa entre 1 y 4 libras de nitrógeno cada 1,000 pies cuadrados por temporada de crecimiento. Cuando se aplica según las instrucciones, este producto proporciona 0.32 libras de nitrógeno cada 1,000 pies cuadrados. Consulte a la oficina local del servicio de extensión para obtener información sobre un programa de aplicación de fertilizante adecuado para su región geográfica.

ESPARCIDORES Y CALIBRACION

Los esparcidores por goteo y los esparcidores giratorios se deben calibrar para lograr la aplicación uniforme de este producto.

Paso único — Opere el equipo a intervalos correspondientes al ancho o a la banda de aspersión del esparcidor. Con esparcidores por goteo, superponga los carriles para asegurar la cobertura total. Cierre las aberturas del esparcidor al girar para cambiar de dirección o cuando se detenga.

IMPORTANTE • Es importante seguir las instrucciones tal como se indican en este producto de calidad para el cuidado del césped, ya que aplicar un exceso de producto puede causar daños a su césped. Los ajustes del esparcidor son aproximados. Puede haber diferencias como consecuencia del estado del esparcidor, la velocidad a la cual se aplica y la irregularidad de la zona tratada. Asegúrese de revisar el esparcidor antes de comenzar y de ajustarlo si es necesario. Consulte el manual del propietario. ***NUNCA LO APLIQUE A MANO.***

Para evitar que queden áreas y líneas sin producto, primero esparza sobre el borde del césped y después complete el centro aplicando en forma paralela y uniforme. Camine a paso regular para aplicar uniformemente el producto en todo el césped.

Cobertura: Para calcular cuánto necesita aplicar, multiplique el largo de su césped por el ancho. Esto equivale a la superficie total en pies cuadrados. Reste todas las áreas que no sean de césped (casa, jardín, etc.). Una bolsa alcanza para cubrir 5,000 pies cuadrados.

AJUSTES DEL ESPARCIDOR

Esparcidores giratorios	
Scotts Edge Guard	6.75
Scotts SpeedyGreen	6.75
Earthway 2050	9.0
Esparcidor por goteo	
Scotts AccuGreen	7.5

INSTRUCCIONES DE CALIBRACION

- Para 3.2 libras de producto cada 1,000 pies cuadrados de césped: verifique que su esparcidor esté regulado en 310 pies cuadrados y ajuste para aplicar la dosis de 1 libra de producto cada 310 pies cuadrados.

ALMACENAMIENTO Y ELIMINACION

WEEDS CONTROLLED/MALEZAS CONTROLADAS

- Annual yellow sweetclover
- Aster
- Austrian fieldcress
- Bedstraw
- Beggarticks
- Betony, Florida
- Bindweed, field
- Bird vetch
- Bitter wintercress
- Bittercress, hairy
- Bitterweed
- Black-eyed Susan
- Black medic
- Black mustard
- Blackseed plantain
- Blessed thistle
- Bloodflower milkweed
- Blue lettuce
- Blue vervain
- Bracted plantain
- Bristly oxtongue
- Broadleaf dock
- Broadleaf plantain
- Broomweed
- Buckhorn
- Buckhorn plantain
- Bulbous buttercup
- Bull thistle
- Bullnettle
- Burclover
- Burdock
- Burning nettle
- Burweed
- Buttercup
- Buttonweed
- Canada thistle
- Carolina geranium
- Carpetweed
- Catchweed bedstraw
- Catnip
- Catsear
- Chickweed, common
- Chickweed, mouseear
- Chicory
- Cinquefoil
- Clover, crimson
- Clover, hop
- Clover, red
- Clover, strawberry
- Clover, sweet
- Clover, white
- Cockle
- Cocklebur
- Common mullein
- Creeping jenny
- Cudweed
- Curly dock
- Daisy, English
- Daisy fleabane
- Daisy, oxeye
- Dandelion
- Dichondra
- Dogfennel
- Dollarweed
- False dandelion
- Falseflax
- False sunflower
- Fiddleneck
- Florida pusley
- Frenchweed
- Galinsoga
- Goldenrod
- Ground ivy
- Gumweed
- Hairy fleabane
- Hawkweed
- Healall
- Heartleaf drymary
- Heath aster
- Hedge bindweed
- Hedge mustard
- Hemp
- Henbit
- Hoary cress
- Hoary plantain
- Hoary vervain
- Horsenettle
- Jimsonweed
- Knawel
- Knotweed
- Kochia
- Lambsquarters
- Lespedeza
- Mallow
- Matchweed
- Mexicanweed
- Milk vetch
- Morningglory
- Mouseear hawkweed
- Mugwort
- Musk thistle
- Mustard
- Narrowleaf plantain
- Narrowleaf vetch
- Nettle
- Orange hawkweed
- Oriental cocklebur
- Oxalis
- Parsley-piert
- Parsnip
- Pearlwort
- Pennycress
- Pepperweed
- Pigweed
- Pineywoods bedstraw
- Plains coreopsis
- Plantain
- Poison ivy
- Poison oak
- Pokeweed
- Poorjoe
- Prairie sunflower
- Prickly lettuce
- Prickly sida
- Prostrate knotweed
- Prostrate pigweed
- Prostrate spurge
- Prostrate vervain
- Puncturevine
- Purslane, common
- Ragweed
- Red sorrel
- Redroot pigweed
- Redstem filaree
- Rough cinquefoil
- Rough fleabane
- Russian pigweed
- Russian thistle
- Scarlet pimpernel
- Scotch thistle
- Sheep sorrel
- Shepherdspurse
- Slender plantain
- Smallflower galinsoga
- Smooth dock
- Smooth pigweed
- Sorrel
- Sowthistle
- Spanishneedles
- Speedwell
- Spiny amaranth
- Spiny cocklebur
- Spiny sowthistle
- Spotted catsear
- Spotted spurge
- Spurweed
- Stinging nettle
- Strawberry, India mock
- Tall nettle
- Tall vervain
- Tansy ragwort
- Tansy mustard
- Tanweed
- Thistle
- Trailing crownvetch
- Tumble mustard
- Tumble pigweed
- Velvetleaf
- Venice mallow
- Virginia buttonweed
- Virginia creeper
- Virginia pepperweed
- Wavyleaf bullthistle
- Western clematis
- Western salsify
- White mustard
- Wild aster
- Wild buckwheat
- Wild carrot
- Wild four-o'clock
- Wild garlic
- Wild geranium
- Wild lettuce
- Wild mustard
- Wild marigold
- Wild onion
- Wild parsnip
- Wild radish
- Wild rape
- Wild strawberry
- Wild sweet potato
- Wild vetch
- Woodsorrel
- Woolly croton
- Woolly morningglory
- Woolly plantain
- Wormseed
- Yarrow
- Yellow rocket
- Yellowflower
- pepperweed and other broadleaf weeds

HOW MUCH TO APPLY

For Kentucky bluegrass, fescues, perennial ryegrass, bermudagrass, bahiagrass, buffalograss, zoysiagrass, St. Augustinegrass and centipedegrass:

Apply 3.2 lb of product per 1,000 sq ft of turf grass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 lb of product per 1,000 sq ft should be used 30 days after the first application.

Do not apply this product to — 'Floratam' St. Augustinegrass in Florida.

The maximum application rate to turf is 1.1 lb 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

A lawn typically utilizes 1 to 4 lb of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides 0.32 lb of nitrogen per 1,000 sq ft. Consult your local Extension service for questions regarding a fertilizer program suitable to your geographical region.

SPREADERS AND CALIBRATION

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

Single Pass — Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.

IMPORTANT • It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult owners' manual. ***NEVER APPLY BY HAND***

To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.

Coverage: To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct any non-lawn areas (house, garden, etc.). One bag covers 5,000 sq ft.

SPREADER SETTINGS

Rotary Spreaders	
Scotts Edge Guard	6.75
Scotts SpeedyGreen	6.75
Earthway 2050	9.0
Drop Spreader	
Scotts AccuGreen	7.5

CALIBRATION INSTRUCTIONS

- For 3.2 lb of product per 1,000 sq ft of turfgrass: Check your spreader on 310 sq ft and adjust to apply at the rate of 1 lb of product per 310 sq ft.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-874-8892 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

IMPORTANTE • Es importante seguir las instrucciones tal como se indican en este producto de calidad para el cuidado del césped, ya que aplicar un exceso de producto puede causar daños a su césped. Los ajustes del esparcidor son aproximados. Puede haber diferencias como consecuencia del estado del esparcidor, la velocidad a la cual se aplica y la irregularidad de la zona tratada. Asegúrese de revisar el esparcidor antes de comenzar y de ajustarlo si es necesario. Consulte el manual del propietario. ***NUNCA LO APLIQUE A MANO.***

Para evitar que queden áreas y líneas sin producto, primero esparza sobre el borde del césped y después complete el centro aplicando en forma paralela y uniforme. Camine a paso regular para aplicar uniformemente el producto en todo el césped.

Cobertura: Para calcular cuánto necesita aplicar, multiplique el largo de su césped por el ancho. Esto equivale a la superficie total en pies cuadrados. Reste todas las áreas que no sean de césped (casa, jardín, etc.). Una bolsa alcanza para cubrir 5,000 pies cuadrados.

AJUSTES DEL ESPARCIDOR

Esparcidores giratorios	
Scotts Edge Guard	6.75
Scotts SpeedyGreen	6.75
Earthway 2050	9.0
Esparcidor por goteo	
Scotts AccuGreen	7.5

INSTRUCCIONES DE CALIBRACION

- Para 3.2 libras de producto cada 1,000 pies cuadrados de césped: verifique que su esparcidor esté regulado en 310 pies cuadrados y ajuste para aplicar la dosis de 1 libra de producto cada 310 pies cuadrados.

ALMACENAMIENTO Y ELIMINACION

ALMACENAMIENTO: Guarde el producto en su envase original y conserve todas las bolsas que contengan producto sin usar en un lugar cerrado, inaccesible a los niños y a los animales domésticos

ELIMINACION DEL ENVASE:

Si está vacío: No vuelva a usar el envase. Tírelo en la basura, o puede ofrecerlo para reciclarlo, si existe esa opción. **Si está parcialmente vacío:** Llame al organismo local que se encarga de los desechos sólidos o llame gratis al 1-800-CLEANUP (1-800-253-2687) para obtener instrucciones sobre su eliminación. Nunca arroje el producto sin usar por ningún desagüe interior ni exterior.

PRECAUCIONES

Es nocivo para los seres humanos y los animales domésticos

PRECAUCION. Es nocivo si se ingiere o si se absorbe por la piel. Causa irritación moderada a los ojos. Evite el contacto con la piel, los ojos o la ropa. Evite respirar el polvo. Cuando utilice este producto, use camisa de manga larga, pantalones largos, calcetines y zapatos. Después de usar este producto, quítese la ropa y lávela por separado antes de volver a usarla y rápidamente lávese muy bien las manos y la piel que estuvo expuesta, con agua y jabón.

RIESGOS PARA EL MEDIO AMBIENTE

El arrastre o el escurrimiento pueden afectar desfavorablemente a la vegetación que no sea el objetivo del tratamiento. No lo aplique directamente al agua. No contamine el agua cuando elimine las aguas de lavado del equipo.

Cuando se aplica según las instrucciones, este producto cumple con las normas para metales aceptadas por la Association of American Plant Food Control Officials.

Se puede obtener información sobre el contenido y las concentraciones de metales de este producto en la siguiente dirección de Internet: http://www.regulatory-info-sb.com

Si usted aplica este producto según las instrucciones y no queda totalmente satisfecho, envíe la prueba de compra de este paquete y el recibo original de la caja registradora a la dirección que se encuentra a continuación. Le devolveremos el valor de compra que pagó.

GARANTIA LIMITADA Y DECLINACION DE RESPONSABILIDAD.

El distribuidor solamente garantiza que la composición química de este producto concuerda con la declaración de ingredientes expresada en la etiqueta, y que el producto es razonablemente adecuado para el uso indicado cuando se aplica según el Modo de empleo.

EL DISTRIBUIDOR NO ESTABLECE NI PROPONE NINGUNA OTRA GARANTIA EXPRESA NI IMPLICITA, INCLUIDA CUALQUIER GARANTIA DE COMERCIABILIDAD O CONVENIENCIA PARA UN PROPOSITO PARTICULAR, LAS QUE SE DECLINAN EXPRESAMENTE. Esta garantía limitada no se extiende al uso del producto de manera incompatible con las instrucciones expresadas en la etiqueta, las advertencias o las precauciones, ni al uso del producto en condiciones anormales, como sequía, lluvias excesivas, tornados, huracanes, etc. Estos factores están fuera del control del distribuidor o del vendedor. Cualquier tipo de daño que surja como consecuencia del incumplimiento de la garantía del distribuidor estará limitado a los daños directos, y no incluirá daños indirectos ni emergentes tales como pérdida de utilidades ni valores, salvo lo dispuesto en otro lugar por la ley.

Los términos de esta garantía limitada y declinación de responsabilidad no se pueden modificar por ningún tipo de declaración ni acuerdo escrito ni verbal. Ningún empleado ni representante del vendedor está autorizado a modificar ni abusar de manera alguna de los términos de esta garantía limitada y declinación de responsabilidad.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at http://www.regulatory-info-sb.com.

If you apply this product as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address below. We will refund the purchase price you paid.

LIMITED WARRANTY AND DISCLAIMER. The distributor warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE DISTRIBUTOR NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the distributor or the seller. Any damages arising from a breach of the distributor's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of the Limited Warranty and Disclaimer in any manner.

Questions or Comments? Call 1-800-874-8892.

Distributed and guaranteed by Chemsico, Division of United Industries Corporation, PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 2217-819-9688

EPA Est. Nos. 9198-OH-1(M), 9198-OH-2(B), 9198-AL-1(A), 49991-TX-1(AZ), 49991-TX-2(BC), 45065-MN-1(BD), 59144-GA-1(BE), 6665-NY-1(BF), 35512-FL-2(BG), 8378-IN-1(AS), 43854-MO-1(BH), 49710-TX-1(AT), 49710-TX-3(AU), 3234-UT-1(AV), 3234-AZ-1(AW), 8660-AL-1(BN), 8660-OH-1(BO), 74764-CA-1(BK), 74764-FL-1(BL), 9404-FL-1(AX), 35497-OR-1(AY)

Circled letter is first letter of lot number. 11-7488 © 2005

Trikote® is a registered trademark of RLC Technologies, L.L.C. The Trikote design is a trademark of RLC Technologies, L.L.C.

FPO

0 13013 54210 8